

CARNITIN

Dietetic additive, aminoacid and vitamin precursor with trace elements.

COMPOSITION:

Carnitine	 30 000	mg
Vitamin D3 (E-671)	 250 000	UI
Vitamin B1	 10 000	mg
Vitamin B2	 4 000	mg
Vitamin B6	 8 000	mg
Vitamin B12	 20	mg
Vitamin C	 20 000	mg
I-Lysin	 30 000	mg
dl-Methionin	 30 000	mg
Copper	 750	mg
Magnesium	 20 000	mg
Manganese	 5 000	mg
Folic acid	 10 000	mg
Nicotinic acid	 50 000	mg
Pantothenic acid	 25 000	mg
Excipient q.s.p.	 1000	g
		-

SPECIES and INDICATIONS:

CARNITIN is indicated for all species, in case of avitaminosis, convalescence and after infectious processes.

PHYSIOLOGIC EFFECTS:

Poultry

- ♦ Improves the muscle formation, reducing the formation of abdominal fat.
- ◆ Reduces the incidence of SDS (Sudden Death Syndrome). The lipid metabolism is improved at heart level.
- ◆ Reduces the cholesterol content in the yolk.
- Improves the hatching rate hatchability.

Swine

- Improves the muscle formation, reducing the formation of back-bone fat.
- Increases boars and sows reproductive performance.

INSTRUCTIONS FOR USE AND DOSE:

Add to the feed at the rate of 2 kg/MT or through the drinking water at the rate of 1 g/L. For boars, add a spoon Carnitin® per animal and day into the individual feeding trough. Add during 8 consecutive days and repeat on a monthly basis.

STORAGE: Keep in a dry and fresh place, protected from light.

PRESENTATION: 1 kg sachets - 25 kg bags

REGISTER No: 26.208 - CAT